

# Effects of solar wind conditions on the plasma wake within a polar crater: preliminary results

M. I. Zimmerman,<sup>1,2,3</sup> W. M. Farrell,<sup>1,3</sup> T. J. Stubbs<sup>1,3,4</sup>

<sup>1</sup> NASA Goddard Space Flight Center, Greenbelt, MD


<sup>2</sup> Oak Ridge Associated Universities, Oak Ridge, TN

<sup>3</sup> NASA Lunar Science Institute, NASA Ames Research Center, Moffett Field, CA

<sup>4</sup> Goddard Earth Sci. and Tech. Center, Univ. MD Baltimore County, Baltimore, MD

michael.i.zimmerman@nasa.gov

**Abstract.** As the solar wind sweeps horizontally past a shadowed lunar crater it simultaneously diffuses toward the surface through an ambipolar process, forming a plasma wake (e.g., Figure 1). Importantly, the resulting electric field structure diverts solar wind protons toward the cold crater floor where they may represent a source of surficial hydrogen. We present a handful of two-dimensional kinetic simulations exploring the range of wake structures and surface particle fluxes possible under various background plasma conditions.



**Figure 1:** Simulated lunar crater wake under typical solar wind conditions for a 500 m deep step-like crater. (*Top*) Net charge concentration [colored lines represent  $\log_{10}(\text{concentration})$  for the respective species; black dashed line is the height at which the Debye length is half the crater depth]. (*Mid.*) Electric field. (*Bot.*) Electric potential.